


## QUIZ #3 – Solutions

### Each problem is worth 5 points

1. The lines are parallel since a vector along each is  $(3, 4, 1)$ .  
 Since  $(1, 0, -2)$  and  $(-1, 2, -5)$  are points on the lines, a second vector in the plane is  $(1, 0, -2) - (-1, 2, -5) = (2, -2, 3)$ .  
 A vector normal to the plane is:  
 $(3, 4, 1) \times (2, -2, 3) = (14, -7, -14)$ , or re-scaling,  $(2, -1, -2)$ .

2. The area of the parallelogram is  
 $|\mathbf{AB} \times \mathbf{AD}| = |(2, 7, 3) \times (1, 1, 1)| = |(4, 1, -5)| = \sqrt{42}$ .



3. If  $(x, y)$  is made to approach  $(1, 1)$  along the straight lines  $y-1 = m(x-1)$ ,

$$\begin{aligned} & \lim_{(x,y) \rightarrow (1,1)} (x^2 - 2x - y^2 + 2y) / (x^2 - 2x + y^2 - 2y + 2) \\ &= \lim_{x \rightarrow 1} ((x-1)^2 - (y-1)^2) / ((x-1)^2 + (y-1)^2) \\ &= \lim_{x \rightarrow 1} ((x-1)^2 - m^2(x-1)^2) / ((x-1)^2 + m^2(x-1)^2) \\ &= \lim_{x \rightarrow 1} (1 - m^2) / (1 + m^2) = (1 - m^2) / (1 + m^2) \end{aligned}$$

Since this result depends on  $m$ , the original limit does not exist.

4.  $\partial f / \partial x = 3x^2/y + \sin(yz/x) + x \cos(yz/x)(-yz/x^2)$   
 $= 3x^2/y + \sin(yz/x) - (yz/x) \cos(yz/x)$

5.  $\nabla f = \exp(x+y+z) \mathbf{i} + \exp(x+y+z) \mathbf{j} + \exp(x+y+z) \mathbf{k} = \exp(x+y+z) (\mathbf{i} + \mathbf{j} + \mathbf{k})$